

AVIAN TUBERCULOSIS in MAMMALS

ANIMAL GROUP AFFECTED	TRANSMISSION	CLINICAL SIGNS	FATAL DISEASE?	TREATMENT	PREVENTION & CONTROL
Primates : Cercopithecidae, rarely Hylobatidae or New World monkeys. Ungulates, ...	Primarily perorally, occasionally cutaneously or aerogenously	Continuous or intermittent diarrhoeas, weight loss, debilitation. In cutaneous infections skin ulcera, in pulmonary infections dyspnoea.	Occasionally (associated with HIV , SIV or SRV infections!)	Usually none, in very valuable animals Cycloserine or Clarithromycin or a combination of Mefloquine, Mefloquine, Mefloquine and Ethambutol	<i>In houses</i> <i>in zoos</i> Indoor housing, strict quarantine and testing programs

<p>Fact sheet compiled by Manfred Brack, formerly German Primate Center, Göttingen / Germany. Alexis Lécu, DVM. Paris Zoo</p>	<p>Last update March 2009</p>
<p>Fact sheet reviewed by W. Rietschel, Wilhelma Zoologischer-Botanischer Garten, Stuttgart, Germany C. Furley, Howletts Zoo, Bekesbourne, United Kingdom</p>	
<p>Susceptible animal groups Primates: Cercopithecidae, rarely Hylobatidae or New World monkeys. Reported also in lemurs (mouse lemurs). Ungulates : deer, bovids, ...</p>	
<p>Causative organism <i>Mycobacterium avium -intracellulare</i> complex with 28 serotypes.</p>	
<p>Zoonotic potential Theoretically yes. (HIV positive humans are most at risk)</p>	
<p>Distribution World – wide.</p>	
<p>Transmission Primarily perorally, occasionally cutaneous or primary pulmonary infections observed. <i>M. avium-intracellulare</i> in contrast to <i>M.tuberculosis / bovis</i> is rarely transmissible between animals but causes environmental (soil, water, structures) contaminations!</p>	
<p>Incubation period</p>	
<p>Clinical symptoms Continuous or intermittent diarrhoeas, weight loss, debilitation, sometimes lymphadenopathy. In cutaneous infections skin ulcerations, in pulmonary infections dyspnoea.</p>	
<p>Post mortem findings Irregular yellowish- white thickening of the intestinal wall (often caecum) due to chronic granulomatous enteritis with infiltration by large, foamy, eosinophilic epithelioid cells or macrophages. Necrosis, fibrosis and calcifications are usually lacking in the enteric lesions. Prominent mesenteric lymphatic channels have been reported. In primary lung infections micro-and macrogranulomas with extensive necrosis were observed.</p>	
<p>Diagnosis Type of morphologic lesions, Acid fast staining of the agents usually present in huge numbers, Tuberculinization, using avian tuberculin, Cultivation of the mycobacteria (slowly growing, thin, transparent, smooth colonies), PCR (misleading in AIDS-patients!).</p>	
<p>Material required for laboratory analysis Faeces, morphological lesions.</p>	

**OIE Reference Laboratory**

- **Dr I. Pavlik**
Veterinary Research Institute
Hudcova 70, 62132 Brno
CZECH (Rep.)
Tel: (420.5) 33.33.16.01 Fax: (420.5) 33.33.12.29
Email: pavlik@vri.cz

Relevant diagnostic laboratories

AFSSA - Mycobactéries
22 rue Pierre Curie,
BP 67, 94703 Maisons-Alfort Cedex
FRANCE
Tel: 01 49.77.13.21 Fax: 01 49.77.13.44 Email: ml.boschiroli@afssa.fr

Nationales Referenzzentrum für Mykobakterien am Forschungszentrum Borstel
D 23845 BORSTEL, Germany
Tel.: 04537 188 213
" " 211
Fax: " " 311

Local medical laboratories.

Treatment

Usually euthanasia, in exceptional cases surgery (granuloma excision) associated with cycloserine or clarithromycin (*M.avium – intracellulare* is resistant to most tuberculostatica!). In Clarithromycin-resistant strains a combination of Mefloquine (40 mg/kg), Mexifloxacin (100 mg/kg) and Ethambutol (100 mg/kg) for 4 weeks has been recommended.

Prevention and control in zoos

Indoor housing of the animals, desinfection of all materials coming in from the environment, strict quarantine and testing program. UV at usual energy/time contact used for drinking water is effective.

Suggested disinfectant for housing facilities**Notification****Guarantees required under EU Legislation****Guarantees required by EAZA Zoos****Measures required under the Animal Disease Surveillance Plan****Measures required for introducing animals from non-approved sources****Measures to be taken in case of disease outbreak or positive laboratory findings****Conditions for restoring disease-free status after an outbreak****Experts who may be consulted**

Frau Dr. S. Rüsck-Gerdes, Nationals Referenzzentrum Borstel.

References

1. Bellinger, D. A., and B. C. Bullock. 1988. Cutaneous *Mycobacterium avium* infection in a cynomolgus monkey. Lab. Anim. Sci. 38: 85 – 86.
2. Bermudez, L. E., P. Kolonoski, M. Petrofsky, M., Wu, C. B. Interlied, and L. S. Young. 2003. Mefloquine, Moxifloxacin, and Ethambutol are a triple – drug alternative to Macrolide – containing regimens for treatment of *Mycobacterium avium* disease. J. Infect Dis. 187: 1977 – 1980.
3. Brack, M. 1987. Agents Transmissible from Simians to Man. Springer, Berlin.
4. Bull, T. J., and D. C. Shanson. 1992. Rapid misdiagnosis by *Mycobacterium avium- intracellulare* masquerading as tuberculosis in PCR/DNA probe tests. Lancet 340: 1360.
5. Frankel, D. H. 1997. Acid fast but slow to grow. Lancet 349: 478.
6. Goodwin, T., C. P. Jerome, and B. C. Bullock. 1988. Unusual lesion morphology and skin test reaction for *Mycobacterium avium* complex in macaques. Lab. Anim. Sci. 38: 20 – 24.
7. Grange, J. M., M. D. Yates, and E. Boughton. 1990. The avian tubercle bacillus and its relatives. J. Appl.

- Bacteriol. 68: 411 – 431.
8. Hatt, J. – M., and F. Guscetti. 1994. A case of mycobacteriosis in a common marmoset (*Callithrix jacchus*). Am. Assoc. Zoo Vet. Annu. Conf. Proc. 1994: 241 – 243.
 9. Hayes, S.L., Sivaganesan, M., White K.M., and Pfaller S.L. 2008. Assessing the effectiveness of low-pressure ultraviolet light for inactivating *Mycobacterium avium* complex (MAC) micro-organisms. Lett Appl Microbiol. 47(5): 386-92.
 10. Heard, D. J., P. E. Ginn, and L. Neuwirth. 1997. *Mycobacterium avium intracellulare* infection in a white – faced saki (*Pithecia pithecia*). J. Zoo Wildl. Med. 28: 185 – 188.
 11. King, N. W. jr. 1993. *Mycobacterium avium – intracellulare* infection. In: Jones, T. C., U. Mohr, and R. D. Hunt. (eds.) Nonhuman Primates I. Monographs on Pathology of Laboratory Animals. Springer, Berlin. Pp. 57 – 63.
 12. Lamberski, N. 1999. Nontuberculous mycobacteria: Potential for zoonosis. In: Fowler, M. E., and R. E. Miller (eds.). Zoo & Wild Animal Medicine. Current Therapy 4. W. B. Saunders, Philadelphia. Pp 146 – 150.
 13. Munson, L., F. J. Luibel, and H. J. van Kruiningen. 1991. Siderophilic bodies associated with hemosiderosis and atypical mycobacterial infection in an island siamang (*Hylobates syndactylus*). J. Med. Primatol. 20: 265 – 270.
 14. Thoen, C. O. 1994. *Mycobacterium avium* infections in animals. Laboratory and clinical aspects of *M. avium* epidemic. II. Forum in microbiology. Res. Microbiol. 145: 173 – 177.
 15. Thorel, M.F., Huchzermeyer, H.F., and Michel, A.L. 2001. *Mycobacterium avium* and *Mycobacterium intracellulare* infection in mammals. Rev Sci Tech. 20(1): 204-18.
 16. Urbain, A. 1950. Deux cas de tuberculose spontanée d'origine aviaire chez un singe africain: Cercopithèque grivet (*Cercopithecus aethiops* L) et chez un singe américain: Oustiti à pinceaux blancs (*Hapale jacchus* L). Bull. Acad. Vét. Fr. 22: 349 – 351.
 17. Weiss, I. K., P. A. Krogstad, C. Botero, L. von Seidlein, and K. Nash. 1995. Fatal *Mycobacterium avium* meningitis after mis – identification of *M. tuberculosis*. Lancet 345: 991 – 992.