

TULAREMIA

ANIMAL GROUP AFFECTED	TRANSMISSION	CLINICAL SIGNS	FATAL DISEASE ?	TREATMENT	PREVENTION & CONTROL
Callitrichids, <i>Saimiri sciureus</i> , <i>Hylobates lar</i> , <i>Macaca</i> spp., <i>Cercopithecus nictitans</i> , <i>Miopithecus talapoin</i> , Lemuridae.	Perorally (food,water), aerogenously, arthropods, contact.	Nonhuman primates: oral ulcers, salivation,nasal/ocular discharges man: pneumonic or typhoid tularemia	<i>Francisella tularensis tularensis</i> in man : 2 – 4%, <i>F.tularensis holoarctica</i> in nonhuman primates fatal	Streptomycin, kanamycin, gentamicin, doxycyclin	<i>in houses:</i> protective clothing during meat processing or skinning of wild animals <i>in zoos:</i> rodent (lagomorph) control

Fact sheet compiled by Manfred Brack, formerly German Primate Center, Göttingen/Germany.	Last update 22.11.2008
Susceptible animal groups <i>F.tularensis tularensis</i> : reservoir hosts: lagomorphs (<i>Sylvilagus</i> spp. , <i>Lepus californicus</i>),sheep; man <i>F.tularensis holoarctica</i> : reservoir hosts: lagomorphs (<i>Lepus timidus</i> , <i>L.europaeus</i>), Cricetidae (<i>Arvicola terrestris</i>), birds. Occasionally: <i>Saguinus oedipus</i> , <i>S. nigricollis</i> , <i>Leontopithecus chrysomelas</i> , <i>Callithrix jacchus</i> , <i>Saimiri sciureus</i> , <i>Hylobates lar</i> , <i>Macaca nukatta</i> , <i>Macaca fascicularis</i> , <i>Cercopitheucus nictitans nictitans</i> , <i>Miopithecus talapoin</i> , <i>Lemur catta</i> , <i>Vaerecia variegata</i> , man	
Causative Organism <i>Francisella tularensis</i> with 4 subspecies: <i>F.tularensis tularensis</i> (Type A), <i>F. tularensis holoarctica</i> (Type B), <i>F.tularensis mediasiatica</i> , <i>F.tularensis novicida</i>).	
Zoonotic potential Yes (10 out of 30 hunters in Hassia/Germany fell ill with tularaemia in 2005!). Altogether 688 human tularaemia cases with at least 2 fatalities in Germany between 1949 and 2006	
Distribution <i>F.tularensis tularensis</i> : North- and South America <i>F.tularensis holoarctica</i> : Europe, Asia, Americas.	
Transmission <i>F. tularensis tularensis</i> : perorally (food, water), aerogenously, arthropods (ticks: <i>Haemaphysalis</i> , <i>Amblyomma</i> , <i>Dermacentor</i> spp.), deer flies (<i>Chrysops discalis</i>), <i>F. tularensis holoarctica</i> : perorally (food, water), tabanids, <i>Culicoides</i> spp.	
Incubation period 2 – 10 days (occasionally more than 1 month)	
Clinical symptoms Callitrichids and <i>M.fascicularis</i> : sudden death, oral ulcers, chronic sinusitis, man: variable from chronic adenopathy to septic infection or pneumonia	
Post mortem findings Ulcerative stomatitis, Lymphadenitis, hepato - and splenomegaly, hepatic and splenic necroses granulomatous pneumonia	
Diagnosis cultivation (aerogenous, blood agar, cysteine heart agar) serology (cave: cross reactions with <i>Brucella</i> spp. and <i>Yersinia enterocolitica</i> !):agglutination, immunofluorescence, ELISA, nested PCR for <i>Fop</i> A gene, RNA sequencing, real time PCR for type A and B differentiation.	
Material required for laboratory analysis blood, pus, necrotic materials (caution: <i>F.tularensis</i> extremely infectious!)	

OIE Reference Laboratory <ul style="list-style-type: none"> Dr T. Mörner National Veterinary Institute, Department of Wildlife 751 89 Uppsala SWEDEN Tel: (46.18) 67.40.00 Fax: (46.18) 67.46.90 Email: torsten.morner@sva.se
Relevant diagnostic laboratories CDC, Atlanta/Georgia, USA CODA, Groeselenberg 99 – 1180, Brussel, Belgium Institut für Mikrobiologie der Bundeswehr 80937 München, Germany Neuherbergstr.11 Tel.: 089 - 3168 - 2312 Fax: 089 - 3168 – 3292 e-mail: InstitutfuerMikrobiologie@bundeswehr.org
Treatment Antibiotics: streptomycin, kanamycin, gentamicin, doxycyclin.
Prevention and control in zoos Proper quarantine management, rodent/ lagomorph control.
Suggested disinfectant for housing facilities
Notification
Guarantees required under EU Legislation
Guarantees required by EAZA Zoos
Measures required under the Animal Disease Surveillance Plan
Measures required for introducing animals from non-approved sources
Measures to be taken in case of disease outbreak or positive laboratory findings
Conditions for restoring disease-free status after an outbreak
Experts who may be consulted: Dr. W. Splettstößer, Dr. E.-J. Finke, Dr. Seibold Institut für Mikrobiologie der Bundeswehr Neuherbergstr. 11 80937 München Tel.: 089 – 3168 – 2312 Fax: 089 – 3168 – 3292 e-mail: wolfsplettstoesser@bundeswehr.org
References <ol style="list-style-type: none"> Brack, M. 1987. Agents Transmissible from Simians to Man. Springer, New York, pp. 171 – 174. Calle, P. P. 1991. An outbreak of tularemia (<i>Francisella tularensis</i>) in zoo primates. Am. Assoc. Zoo Vet. Annu Conf Proc. 1991: 218. Dedek, J., H. Loepelmann, und R. Kokles. 1990. Ergebnisse serologischer Untersuchungen auf ausgewählte Infektionen bei Feldhasen. Verh. Ber. Erkr. Zootiere 32 : 199 – 204. Doyle, L., J. Markovits, and J. Roberts. 1988. Tularemia (<i>Francisella tularensis</i>) in a squirrel monkey (<i>Saimiri sciureus</i>). Lab. Anim. Sci. 38 : 491 – 492. Enderlin, G., L. Morales, R. F. Jacobs, and J. T. Cross. 1994. Streptomycin and alternative agents for the treatment of tularemia : review of the literature. Clin. Infect. Dis. 19 : 42 – 47. Fulop, M., D. Leslie, and R. Titball. 1996. A rapid, highly sensitive method for the detection of <i>Francisella tularensis</i> in clinical samples using the polymerase chain reactions. Am. J. Trop. Med. Hyg. 54 : 364 – 366. Grunow, R., and H. S. Priebe. 2007. Tularämie – Zum Vorkommen in Deutschland. Epidemiol. Bull. RKI (7), 51 – 56.

8. Hoelzle, L. E., L. Corboz, P. Ossent, and M. M. Wittenbrink. 2004. Tularemia in a captive golden-headed lion tamarin (*Leontopithecus chrysomelas*) in Switzerland. *Vet. Rec.* 155: 60 – 61.
9. Kugeler, K. J., R. Pappert, Y. Zhou, and J. M. Petersen. 2006. Real-time PCR for *Francisella tularensis* Types A and B. *Emerg. Infect. Dis.* 12 : 1799 – 1801.
10. Leong, K. M., S. P. Terrell, and A. Savage. 2004. Causes of mortality in captive cotton-tailed tamarins (*Saguinus oedipus*). *Zoo Biol.* 23 : 127 – 137.
11. Line, A.S., J. E. Markovits, and R. P. Tarara. 1991. Nonhuman primate tularemia: a review of four cases. *Am. Assoc. Lab. Anim. Bull.* 30 (4): 18.
12. Maetz-Rensing, K., A. Floto, A. Schrod, T. Becker, E. J. Finke, E. Seibold, W. D. Spletstoesser, and F.-J. Kaup (2007) : Epizootic of tularemia in an outdoor housed group of cynomolgus monkeys (*Macaca fascicularis*). *Vet. Pathol.* 44 : 327 – 334.
13. Mignani, E., F. Palmieri, M. Fontana, and S. Marigo. 1988. Italian epidemic of waterborne tularemia. *Lancet* II : 1423.
14. Morrow, G. W., P. A. Day – Lollini, P. A. Murphy – Hackley, J. R. Blum, and T. E. Hamm, jr. . 1990. Tularemia in a laboratory maintained squirrel monkey. *Lab. Anim. Sci.* 40 : 568.
15. Posthaus, H., M. Welle, T. Mörner, J. Nicolet, and P. Kuhnert. 1998. Tularemia in a common marmoset (*Callithrix jacchus*) diagnosed by 16 S rRNA sequencing. *Vet. Microbiol.* 61 : 145 – 150.
16. Preiksaitis, J. K., G. J. Crawshaw, G. S. P. Nayar, and H. G. Stiver. 1979. Human tularemia at an urban zoo. *Can. Med. Assoc. J.* 121 : 1097 – 1099.
17. Waggle, K. S., P. A. Day – Lollini, P. A. Murphy – Hackley, J. R. Blum, and G. W. Morrow. 1997. Diagnostic exercise : Illness, cutaneous hemorrhage, and death in two squirrel monkeys (*Saimiri sciureus*). *Lab. Anim. Sci.* 47 : 647 – 649.