

## BOVINE BABESIOSIS

ANIMAL GROUP AFFECTED	TRANSMISSION	CLINICAL SIGNS	FATAL DISEASE ?	TREATMENT	PREVENTION & CONTROL
Bovine	Tick-borne	Fever, lethargy, diarrhoea, anaemia, haemoglobinuria icterus hyperaesthesia, convulsions, paralysis	Yes	Diminazene or imidocarb	<i>In houses</i> Tick control  <i>in zoos</i> Tick control

<b>Fact sheet compiled by</b> J. Brandt, Royal Zoological Society of Antwerp, Belgium	<b>Last update</b> February 2009
<b>Fact sheet reviewed by</b> D. Geysen, Animal Health, Institute of Tropical Medicine, Antwerp, Belgium F. Vercammen, Royal Zoological Society of Antwerp, Belgium	
<b>Susceptible animal groups</b> The most important species in cattle ( <i>Babesia bovis</i> , <i>Babesia bigemina</i> and <i>Babesia divergens</i> ) are in general more pathogenic in adult animals than in calves. Infection occurs equally in buffalo, bison and deer. <i>B. divergens</i> also in rodents and primates.	
<b>Causative organism</b> Several species belonging to the phylum of the Apicomplexa, order Piroplasmida, family Babesiidae; Pathogenic species are <i>B. bigemina</i> , <i>B. bovis</i> and <i>B. divergens</i> . Babesiosis has been dubbed piroplasmosis, tick fever, Texas fever, redwater. Other species, i.a. <i>Babesia major</i> , <i>Babesia occultans</i> are considered to be less or non pathogenic.	
<b>Zoonotic potential</b> <i>B. divergens</i> or <i>B. divergens</i> -like parasites (and the rodent parasite <i>Babesia microti</i> in the USA) are considered as emerging zoonoses often studied in association with other tick-borne infections such as Lyme disease and human granulocytic anaplasmosis/ehrlichiosis or in immunocompromised or splenectomised patients.	
<b>Distribution</b> <i>B. bigemina</i> and <i>B. bovis</i> : South and Central America, Europe, Africa, Middle East, Central Asia, Australia. <i>B. divergens</i> : common <i>Babesia</i> sp. of cattle in Europe. <i>B. major</i> : North Africa and Europe.	
<b>Transmission</b> Ticks of the genera <i>Boophilus</i> , <i>Rhipicephalus</i> , <i>Haemaphysalis</i> or <i>Ixodes</i> with transovarial transmission as a rule. Specificity for the definite host can be very strict and is decisive for the distribution; e.g. <i>B. bovis</i> and <i>B. bigemina</i> are mainly transmitted by the one-host <i>Boophilus</i> spp. where <i>Boophilus decoloratus</i> transmits <i>B. bigemina</i> but not <i>B. bovis</i> , for the latter <i>Boophilus microplus</i> is the main vector. <i>B. divergens</i> is transmitted by the 3-host tick, <i>Ixodes ricinus</i> . <i>Haemaphysalis punctata</i> is an important host of <i>B. major</i> . but it occurs equally in <i>Boophilus</i> sp. and in <i>Ixodes</i> sp. Mechanical transmission is possible.	
<b>Incubation period</b> In general one to two weeks p.i. but this and the course of the infection depend on many factors: inoculum, age, breed and physiological status of the intermediate host. Adults are more susceptible than calves, European breeds and Zebu (except for <i>B. bovis</i> ) more susceptible than African breeds, perinatal suppression and concomitant infections (e.g. <i>Anaplasma (Ehrlichia) phagocytophila</i> and <i>B. divergens</i> ) are known to occur.	
<b>Clinical symptoms</b> Fever, lethargy, diarrhoea, haemolytic anaemia, tachycardia, haemoglobinuria and icterus are common features of acute infections. Chronic infections often without apparent haemoglobinuria. Acute <i>B. bovis</i> often develops in fatal cerebral babesiosis with hyperaesthesia, convulsions and paralysis due to aggregation of red blood cells in the cerebral capillaries and extravascular, following endothelial damage. No or little pathogenicity contributed to <i>B. major</i> . Recovered animals become carriers, without apparent clinical	



symptoms, but with possibility to relapse under stress conditions. They also remain infective.

**Post mortem findings**

Icteric appearance of the tissues, splenomegaly, congested liver with distended gall bladder, kidneys congested and haemorrhagic, subcutaneous and pulmonary oedema, sero-haemorrhagic fluid in all cavities, urinary bladder with reddish or brownish urine (not in chronic babesiosis). In cases of *B.bovis* with nervous symptoms, petechiae and oedematous congestion of the brain. Petechiae and ecchymosis in the epi- and myocard and the kidneys.

**Diagnosis**

Direct diagnosis: piriform merozoites in blood smears after staining (Giemsa after fixation) preferably from capillary blood (*B.bovis*) or at necropsy, in brain smears (*B.bovis*) where clumps of infected erythrocytes can be observed in microthrombi.

*B.bigemina*: relatively large merozoites round (2-3 µm) or irregular elongated shape (up to 5 µm), also in blood from the general circulation. Paired forms often in acute angle.

*B.bovis*: parasitaemias usually very low. Mostly annular vacuolised forms. Paired forms often in obtuse angle. Piroplasms measure 1.5-2 µm and are in the centre of the cells.

*B.divergens*: small merozoites ring-shaped or piriform, when paired in an obtuse angle 1.5-2 µm, often at the border of infected cells.

*B.major*: resembles *B.bovis*, centrally positioned, annular forms measure 1.8 µm, elongated (pear-) shaped forms 2.6-3.7 µm.

PCR and antibody detection assays. Immunofluorescent antibody test (IFAT) and slide-enzyme-linked immunosorbent assay (SELISA) are satisfactory but cross reactions between *Babesia* spp. do not allow speciation. Antigen slides are prepared from controlled infections in splenectomised bovines or from *Meriones* sp. (*B.divergens*).

**Material required for laboratory analysis**

Thin blood smears or EDTA-anticoagulated blood. When *B.bovis* is suspected preferably capillary blood has to be taken and for cerebral *B.bovis*, smears of brain capillaries.

Parasitaemias in latent infections are often too low to be detected, thick smears sometimes prepared from RBC just underneath the buffy coat and repeated sampling can improve detection.

**OIE Reference Laboratory**

- **Prof. Ikuo Igarashi**  
National Research Center for Protozoan Diseases, Obihiro University of Agriculture and Veterinary Medicine  
Inada-cho Nishi 2-13, Obihiro, Hokkaido 080-8555  
JAPAN  
Tel: (81.155) 49.56.42 Fax: (81.155) 49.56.43  
Email: [igarcpmi@obihiro.ac.jp](mailto:igarcpmi@obihiro.ac.jp)

**Relevant diagnostic laboratories**

CODA, Groeselenberg 99, 1180 Brussel, Belgium

**Treatment**

Diamidine derivatives are well tolerated and have good activity (1) Diminazene at 3.5 mg/kg b.w. i.m. curative and (2) Imidocarb at 1.2 mg/kg b.w. i.m. or s.c. curative or 3 mg/kg b.w. prophylactic (for 3 to 6 weeks).

**Prevention and control in zoos**

Tick control by acaricidal treatment or vaccination in domestic animals might be superfluous in a supposedly tick free environment of the zoo, but specific attention to tick prevention in imported animals is recommended.

**Suggested disinfectant for housing facilities****Notification****Guarantees required under EU Legislation****Guarantees required by EAZA Zoos****Measures required under the Animal Disease Surveillance Plan****Measures required for introducing animals from non-approved sources****Measures to be taken in case of disease outbreak or positive laboratory findings****Conditions for restoring disease-free status after an outbreak**

**Contacts for further information****References**

1. Bock RE, Kingston TG, Devos AJ (1999) Effect of breed of cattle on transmission rate and innate resistance to infection with *Babesia bovis* and *Babesia bigemina* transmitted by *Boophilus microplus*. *Austr Vet J* 77: 461-464.
2. Brown CDG, Hunter AG, Luckins AG. Bovine babesiosis. In: Sewell MMH, Brocklesby DW (eds.). *Handbook on Animal Diseases in the Tropics*, Chapter: Diseases caused by protozoa. Baillière Tindall, 1990, 161-170.
3. Brunhansen H, Christensson DA, Hardeng F, Gronstol H (1997) Experimental infection with *Ehrlichia phagocytophila* and *Babesia divergens* in cattle. *J Vet Med* 44: 235-243.
4. De Vos AJ, Potgieter FT. Bovine babesiosis. In: *Infectious Diseases of Livestock*. Coetzer JAW, Thomson GR, Tustin RC (eds.) Oxford University Press, 1994, 277-294.
5. Gray J, von Stedingk L.V, Granstrom M (2002) Zoonotic babesiosis. *Int J Med Microbiol* 291: 108-111.
6. L'Hostis M, Seegers H (2002) Tick-borne parasitic diseases in cattle: current knowledge and prospective risk analysis related to the ongoing evolution in French cattle farming systems. *Vet Res* 33: 599-611.
7. Kanyari PWN, Kagira J (2000) The role of parasitic diseases as causes of mortality in cattle in a high potential area of central Kenya: a quantitative analysis. *Onderstepoort J Vet Res* 67: 157-161.
8. Kocan AA., Waldrup KA. Piroplasms (*Theileria* spp., *Cytauxzoon* spp., and *Babesia* spp.). In: Samuel WM, Pybus MJ, Kocan AA (eds.). *Parasitic Diseases of Wild Mammals*, Chapter 18: Blood-Inhabiting Protozoans. Iowa State University Press, 2001, 524-536.
9. Schettlers TPM, Kleuskens J, Scholtes N, Gorenflot A (1998) Parasite localization and dissemination in the *Babesia*- infected host. *Ann Trop Med Parasitol* 92: 513-519.
10. Sevinc F, Sevinc M, Birdane FM, Altinoz F (2001) Prevalence of *Babesia bigemina* in cattle. *Rev Med Vet* 152: 395-398.
11. Taylor M (2000) Protozoal disease in cattle and sheep. *In Practice* 22: 604-617.